

# Mr. Kent's Organic Chemistry Unit Notes


## I Basic Concepts

A. Organic Chemistry-The study of \_\_\_\_\_ containing compounds.


1. They occur extensively in nature because all living things are made of \_\_\_\_\_ containing compounds.

## II Bonding

A. \_\_\_\_\_ can form \_\_\_\_\_ covalent bonds (\_\_\_\_\_ hybrids consisting of 1 \_\_\_\_\_ orbital and 3 \_\_\_\_\_ orbitals)




When hybridized the 4 different orbitals are combined and create 4 equal energy orbitals.



B. \_\_\_\_\_ will not only combine with other atoms, but will also bond with other \_\_\_\_\_ atoms making for very large numbers of molecules.

C. Organic compounds are more numerous than \_\_\_\_\_ compounds.

D. A major source of organic compounds is \_\_\_\_\_.  
Which came from \_\_\_\_\_ many years ago.




### III Characteristics

1. Generally, \_\_\_\_\_ molecules (SNAP)
2. \_\_\_\_\_ dissolves in water  
Exceptions \_\_\_\_\_ & \_\_\_\_\_
3. Non-\_\_\_\_\_ -generally do not conduct electricity in water, except \_\_\_\_\_
4. Melting Points are \_\_\_\_\_ due to \_\_\_\_\_ intermolecular forces
5. Chemical Reactions are \_\_\_\_\_ compared to inorganic reactions

### IV Some Organic Compounds form \_\_\_\_\_.

1. Compounds with the same \_\_\_\_\_ but different \_\_\_\_\_
2. We draw structural formulas to avoid confusion
3. Example- $C_3H_8O$



4. As the number of carbon atoms increase, the number of Isomers will \_\_\_\_\_.

#### V More bonding

1. Carbon atoms that are bonded together by sharing \_\_\_\_\_ electrons form a \_\_\_\_\_ bond
2. Carbon atoms that are bonded together by sharing \_\_\_\_\_ electrons form a \_\_\_\_\_ bond
3. Carbon atoms that are bonded together by sharing \_\_\_\_\_ electrons form a \_\_\_\_\_ bond
4. \_\_\_\_\_ compounds contain ALL SINGLE BONDS
5. \_\_\_\_\_ compounds contain NOT ALL SINGLE BONDS, but have double or triple bonds.

#### VI. Types of Organic Compounds


##### A. HYDROCARBONS

1. Contain only \_\_\_\_\_ and \_\_\_\_\_ atoms.
2. Homologous Series (see table \_\_\_\_\_)- each sample of a homologous series differs by the previous by \_\_\_\_\_ carbon and \_\_\_\_\_ hydrogen atoms
3. Boiling Points- As the number of carbon atoms increases the boiling points \_\_\_\_\_ due to \_\_\_\_\_ intermolecular forces.

## ALKANES- Saturated hydrocarbons


Nomenclature (Naming Rules)- end with “-ane”

General Formula




# Carbons	prefix	Molecular formula	Name
1	Meth	CH <sub>4</sub>	
2	Eth	C <sub>2</sub> H <sub>6</sub>	
3	Prop	C <sub>3</sub> H <sub>8</sub>	
4	But	C <sub>4</sub> H <sub>10</sub>	
5	Penta	C <sub>5</sub> H <sub>12</sub>	
6	Hexa	C <sub>6</sub> H <sub>14</sub>	
7	Hept	C <sub>7</sub> H <sub>16</sub>	
8	Oct	C <sub>8</sub> H <sub>18</sub>	
9	Non	C <sub>9</sub> H <sub>20</sub>	
10	Deca	C <sub>10</sub> H <sub>22</sub>	

The first 3 alkanes have no isomers (they can only be drawn 1 way).




## Isomers of Butane C<sub>4</sub>H<sub>10</sub>


**Rule #1**-Name the longest continuous chain of carbon atoms (with group), and end it with -ane.




Rule #2- Remaining side chains will be given the ending -yl.



### Isomers of Pentane C<sub>5</sub>H<sub>12</sub>



Rule #3- Multiple side chains will use prefixes 2 is di-, 3 is tri-, 4 is tetra- and so on.







## Isomers of Hexane C<sub>6</sub>H<sub>14</sub>

$  \begin{array}{cccccc}  & \text{H} & \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \\  &   &   &   &   &   &   \\  \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\  &   &   &   &   &   &   \\  & \text{H} & \text{H} & \text{H} & \text{H} & \text{H} & \text{H}  \end{array}  $	
$  \begin{array}{cccccc}  & & \text{H} & & & & \\  & &   & & & & \\  & & \text{H} - \text{C} - \text{H} & & & & \\  & &   & & & & \\  & \text{H} & & \text{H} & \text{H} & \text{H} & \\  &   & &   &   &   & \\  \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\  &   & &   &   &   & \\  & \text{H} & & \text{H} & \text{H} & \text{H} &   \end{array}  $	
$  \begin{array}{cccccc}  & & & \text{H} & & & \\  & & &   & & & \\  & & & \text{H} - \text{C} - \text{H} & & & \\  & & &   & & & \\  & \text{H} & \text{H} & & \text{H} & \text{H} & \\  &   &   & &   &   & \\  \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\  &   &   & &   &   & \\  & \text{H} & \text{H} & & \text{H} & \text{H} &   \end{array}  $	

**Rule #4-** Give the lowest number location for each side chain if there is another possible location it could be located. Go back and fix the last 2 isomers





# Isomers of Heptane $C_7H_{16}$

$  \begin{array}{cccccccc}  & H & H & H & H & H & H & H \\  &   &   &   &   &   &   &   \\  H & - C & - C & - C & - C & - C & - C & - H \\  &   &   &   &   &   &   &   \\  & H & H & H & H & H & H & H  \end{array}  $	
$  \begin{array}{ccccccc}  & & H & & & & \\  & &   & & & & \\  & & H-C-H & & & & \\  & &   & & H & H & H & H \\  & H & - C & - C & - C & - C & - C & - H \\  &   &   &   &   &   &   &   \\  & H & H & H & H & H & H & H  \end{array}  $	
$  \begin{array}{ccccccc}  & & & H & & & \\  & & &   & & & \\  & & & H-C-H & & & \\  & H & H &   & H & H & H \\  &   &   &   &   &   &   \\  H & - C & - C & - C & - C & - C & - H \\  &   &   &   &   &   &   \\  & H & H & H & H & H & H  \end{array}  $	
$  \begin{array}{ccccccc}  & & H & & & & \\  & &   & & & & \\  & & H-C-H & & & & \\  & H &   & H & H & H & \\  &   &   &   &   &   & \\  H & - C & - C & - C & - C & - C & - H \\  &   &   &   &   &   & \\  & H & H-C-H & H & H & H & \\  & &   & & & & \\  & & H & & & &   \end{array}  $	
$  \begin{array}{ccccccc}  & & H & & & & \\  & &   & & & & \\  & & H-C-H & & & & \\  & H &   & H & H & H & \\  &   &   &   &   &   & \\  H & - C & - C & - C & - C & - C & - H \\  &   &   &   &   &   & \\  & H & H & H-C-H & H & H & \\  & & &   & & & \\  & & & H & & &   \end{array}  $	

















**Rule #5-** When there are 2 different side chains name them in alphabetical order using the carbon prefix (meth, eth..).





## 2. Alkenes-

A. Are \_\_\_\_\_ (not all single bonds) hydrocarbons (hydrogen and carbon only) containing 1 carbon to carbon double bond "C=C".



B. The rule for naming is they all end with "-ene".

C. The general formula is  $C_nH_{2n}$ ,  $n$  is the number of carbons is used to determine the number of hydrogen atoms. Example  $n=5$ , so  $H=(2(5))=10$




What is the first member of the Alkene homologous series?

Name \_\_\_\_\_ Formula \_\_\_\_\_

**Rule #1-**Name the longest chain with the double bond. It is not always the longest chain of carbons. End it with "-ene".

C <sub>2</sub> H <sub>4</sub>		
C <sub>3</sub> H <sub>6</sub>		

**Rule #2-**When necessary use the lowest number to give the location of double bond in the longest chain.

C <sub>4</sub> H <sub>8</sub>		
C <sub>4</sub> H <sub>8</sub>		
C <sub>5</sub> H <sub>10</sub>		





C. **Alkynes**- Are \_\_\_\_\_ (not all single bonds) hydrocarbons (hydrogen and carbon only) containing 1 carbon to carbon double bond " $C\equiv C$ ".

The rule for naming is they all end with "-yne".

The general formula is  $C_nH_{2n-2}$ , n is the number of carbons is used to determine the number of hydrogen atoms. Example  $n=5$ , so  $H=(2(5)-2)=8$

**Rule #1**-Name the longest chain with the triple bond. It is not always the longest chain of carbons. End it with "-yne".

$C_2H_2$	$H-C\equiv C-H$	
$C_3H_4$	$  \begin{array}{c}  H \\    \\  H-C\equiv C-C-H \\    \\  H  \end{array}  $	

**Rule #2**-When necessary use the **lowest** number to give the location of triple bond in the longest chain.

$C_3H_4$	$  \begin{array}{c}  H \\    \\  H-C\equiv C-C-H \\    \\  H  \end{array}  $	
$C_4H_6$	$  \begin{array}{c}  H \quad H \\    \quad   \\  H-C\equiv C-C-C-H \\    \quad   \\  H \quad H  \end{array}  $	

C <sub>4</sub> H <sub>6</sub>	$  \begin{array}{c}  \text{H} \qquad \qquad \text{H} \\    \qquad \qquad   \\  \text{H}-\text{C}-\text{C}\equiv\text{C}-\text{C}-\text{H} \\    \qquad \qquad   \\  \text{H} \qquad \qquad \text{H}  \end{array}  $	
C <sub>5</sub> H <sub>8</sub>	$  \begin{array}{c}  \qquad \qquad \text{H} \quad \text{H} \quad \text{H} \\  \qquad \qquad   \quad   \quad   \\  \text{H}-\text{C}\equiv\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\  \qquad \qquad   \quad   \quad   \\  \qquad \qquad \text{H} \quad \text{H} \quad \text{H}  \end{array}  $	
C <sub>5</sub> H <sub>8</sub>	$  \begin{array}{c}  \qquad \qquad \text{H} \quad \text{H} \\  \qquad \qquad   \quad   \\  \text{H}-\text{C}-\text{C}\equiv\text{C}-\text{C}-\text{C}-\text{H} \\    \qquad \qquad   \quad   \\  \text{H} \qquad \qquad \text{H} \quad \text{H}  \end{array}  $	


## 4. Benzenes-

The rule for naming is they all end with "benzene".


The general formula is  $C_nH_{2n-6}$ ,  $n$  is the number of carbons is used to determine the number of hydrogen atoms. Example  $n=8$ , so  $H=(2(8)-6)=10$

**Rule #1**-Name the benzene ring. End it with "benzene".



**Rule #2**-When necessary use the lowest number to give the locations of side chains that come off the benzene.



$C_8H_{10}$



$C_9H_{12}$



## 5. Other Hydrocarbons




**A. Dienes-** Organic compound containing 2 separate carbon to carbon double bonds C=C

The rule for naming is they all end with "-diene".

The general formula is  $C_nH_{2n-2}$ , n is the number of carbons is used to determine the number of hydrogen atoms. Example n=4, so  $H=(2(4)-2)=6$

**Rule #1-**Name the longest chain that contains both double bonds. End it with "diene".

**Rule #2-**When necessary use the ***lowest*** numbers to give the locations of both double bonds.

$C_4H_6$		
$C_5H_8$		
$C_5H_8$		





B. **Trienes**- Organic compound containing 3 separate carbon to carbon double bonds C=C

The rule for naming is they all end with "-triene".

The general formula is  $C_nH_{2n-4}$ , n is the number of carbons is used to determine the number of hydrogen atoms. Example n=6, so  $H=(2(6)-4)=8$

**Rule #1**-Name the longest chain that contains both double bonds. End it with "triene".





**Rule #2**-When necessary use the lowest numbers to give the locations of both double bonds.

C <sub>6</sub> H <sub>8</sub>		
C <sub>6</sub> H <sub>8</sub>		
C <sub>6</sub> H <sub>8</sub>		
C <sub>7</sub> H <sub>10</sub>		

C. **Cyclic compounds**-Hydrocarbons in which the 2 ends of the chain are attached at the ends to form a ring

The rule for naming is they all start with "cyclo-".

**Rule #1**-Name the carbon ring. Start it with "cyclo".

$C_3H_6$		
$C_4H_8$		
$C_5H_{10}$		
$C_6H_{12}$		

## B. Other Organic Compounds

1. Functional Groups-the atom or group of atoms that characterizes the structure of a family of organic compounds and determines the properties

### 2. Halides

A. An organic compound containing one or more halogen atoms.

B. **Nomenclature**-Name the halogen like you would any other substituent group.

Chlorine-->Chloro      Bromine-->Bromo

Iodine-->Iodo          Fluorine-->Fluoro

### C. Examples

Name	Structure
	$\begin{array}{c} \text{F} \\   \\ \text{Cl}-\text{C}-\text{Cl} \\   \\ \text{F} \end{array}$
	$\begin{array}{c} \text{Cl} \\   \\ \text{Cl}-\text{C}-\text{H} \\   \\ \text{Cl} \end{array}$

3. Alcohols- are compounds in which one or more hydrogen atoms in an alkane have been replaced by a hydroxyl (-OH) group.


OH- is hydroxide, but in organic chemistry side chains end with “-yl”.





## Types of Alcohols

Monohydroxy Alcohols- contains 1 -OH group



**Primary(1°)**- the C-OH is attached to one other carbon (on the end)

**Secondary(2°)**- the C-OH is attached to two other carbons

**Tertiary(3°)**- the C-OH is attached to three other carbons

Primary Example



Secondary






Tertiary Example




Dihydroxy Alcohols- contains 2 –OH group (ends with -\_\_\_\_\_)



Trihydroxy Alcohols- contains 3 –OH group (ends with -\_\_\_\_\_)




4. Organic Acids- An organic acid is an organic compound with acidic properties. The most common organic acids are the carboxylic acids whose acidity is associated with their carboxyl group -COOH.





**Nomenclature**-carboxylic acids have an -oic acid suffix (e.g. octadecanoic acid)

Name	Expanded Structure
Name	Condensed Structure
	HCOOH
	CH <sub>3</sub> COOH
	CH <sub>3</sub> CH <sub>2</sub> COOH
	C <sub>4</sub> H <sub>9</sub> COOH
	C <sub>8</sub> H <sub>17</sub> COOH



5. Aldehydes-A compound with a carbon atom which is bonded to a hydrogen atom and double-bonded to an oxygen atom. This is found on the end of the chain, ketones have this carbon in the middle (no H either).

**Nomenclature**-The name is formed by changing the suffix -e of the parent alkane to -al, so that HCHO is named methanal, and CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CHO is named butanal.


Expanded Structure	Name	Expanded Structure	Name
			

## Synthesis




A primary alcohol reacts with an oxidizing agent to make an aldehyde



6. Ketones- A compound with a carbonyl group (O=C) linked to two other carbon atoms in the chemical compound. The C=O is found somewhere in the middle of the chain (not the end). Three carbons are needed to form a ketone.



**Nomenclature**- changing the suffix -e of the parent alkane to “-one”


Name	Expanded Structures	Name	Condensed Structures
			CH <sub>3</sub> COCH <sub>3</sub>
			CH <sub>3</sub> COCH <sub>2</sub> CH <sub>3</sub>
			CH <sub>3</sub> COCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>
			CH <sub>3</sub> CH <sub>2</sub> COCH <sub>2</sub> CH <sub>3</sub>
			CH <sub>3</sub> COCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>

## Synthesis




Ketones can be created by oxidation of secondary alcohols. The process requires a strong oxidizing agent.



7. **Ethers**-a chemical compound where an oxygen atom connects two substituted groups




**Nomenclature**-Ethers can be named by naming each of the two carbon groups as a separate word followed by a space and the word ether.

Name	Expanded Structure
	
	
	
	$\text{CH}_3\text{-CH}_2\text{-O-CH}_2\text{-CH}_2\text{-CH}_3$
	$\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O-CH}_2\text{-CH}_2\text{-CH}_3$

## Synthesis

Primary alcohols react to produce an ether and water




## 8. Polymers-composed of many repeating units called monomers



Natural Polymers	Synthetic Polymers



## 9. Esters

**Nomenclature**-Esters ( $R-CO-O-R'$ ) are named as alkyl derivatives of carboxylic acids. The alkyl ( $R'$ ) group is named first (the carbon chain from the alcohol). The  $R-CO-O$  part is then named as a separate word based on the carboxylic acid name, with the ending changed from *-oic acid* to *"-oate"*.






**"Alcoholyl acidoate"**

NAMES	Expanded STRUCTURE
	
	


	
	
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{COOCH}_3$
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{COOCH}_3$
	$\text{CH}_3\text{CH}_2\text{COOCH}_3$

10. Amines-In amines, the hydrogen atoms in the ammonia have been replaced one at a time by hydrocarbon groups. On this page, we are only looking at cases where the hydrocarbon groups are simple alkyl groups.




**11. Amino Acids** are the chemical units or "building blocks" of the body that make up proteins. Contain both an amine group (-NH<sub>2</sub>) and the carboxyl group (-COOH)



**12. Amides-** (R-CO-NH<sub>2</sub>) take the suffix "-amide". There is no prefix form, and no location number is required since they always terminate a carbon chain, e.g. CH<sub>3</sub>CONH<sub>2</sub> is named ethanamide.






## VII. Organic Reactions


**A. Substitution reaction**-A functional group in a particular chemical compound is replaced by another

This occurs with a saturated structure.


### Saturated, Substitution





**B. Addition reaction**-A reaction where a pi bond (either a double or triple bond) is replaced with the creation of two new covalent bonds. (for this course you only need to know the term addition and how it works)



### Hydrogenization




### Halogenation

-bromine or chlorine


### C. Fermentation

The enzymes in yeast break down sugar (glucose) into alcohol (ethanol) and carbon dioxide gas.




D. **ESTERIFICATION**-is the general name for a chemical reaction in which two chemicals (typically an acid and an alcohol) form an ester as the reaction product (water).

**Synthesis**-When a carboxylic acid and an alcohol react, a water molecule is removed, and an ester molecule is formed.



E. **Saponification** is commonly used to refer to the reaction of a **fat** (ester) with a metallic alkali (**base**) or oil to form **soap** and **glycerol**.



F. **Oxidation-(combustion)**- Complete combustion (given sufficient oxygen) of any hydrocarbon produces carbon dioxide and water.



Incomplete Combustion-not enough O<sub>2</sub>



## G. Polymerization

1. Joining of \_\_\_\_\_
2. Condensation Polymerization (evolves \_\_\_\_\_)



Monomers

Dimer

Dimer is 2 monomers combined


Trimer

Polymer


The prerequisite is 2 coupling groups on each end

### 3. Addition Polymerization

A double or triple bond is broken and a monomer adds to the chain




**Note:**  $n$  and  $n$  are very large numbers equal to about 2000.



## H. Petroleum Reactions

1. **Fractional Distillation**- The various components of crude oil have different sizes, weights and boiling temperatures; so, the first step is to separate these components. Because they have different boiling temperatures, they can be separated easily by a process called fractional distillation.



2. **Cracking**-Large hydrocarbon are broken into smaller ones, yielding more valuable fractions (gasoline, fuel oils) using a catalyst.

